

ON ABSOLUTE VALUES OF \mathcal{Q}_K FUNCTIONS

GUANLONG BAO, ZENGJIAN LOU, RUISHEN QIAN, AND HASI WULAN

ABSTRACT. In this paper, the effect of absolute values on the behavior of functions f in the spaces \mathcal{Q}_K is investigated. It is clear that $f \in \mathcal{Q}_K(\partial\mathbb{D}) \Rightarrow |f| \in \mathcal{Q}_K(\partial\mathbb{D})$, but the converse is not always true. For f in the Hardy space H^2 , we give a condition involving the modulus of the function only, such that this condition together with $|f| \in \mathcal{Q}_K(\partial\mathbb{D})$ is equivalent to $f \in \mathcal{Q}_K$. As an application, a new criterion for inner-outer factorisation of \mathcal{Q}_K spaces is given. These results are also new for \mathcal{Q}_p spaces.

1. INTRODUCTION

Denote by $\partial\mathbb{D}$ the boundary of the unit disk \mathbb{D} in the complex plane \mathbb{C} . Let $H(\mathbb{D})$ be the space of functions analytic in \mathbb{D} . Throughout this paper, we assume that $K : [0, \infty) \rightarrow [0, \infty)$ is a right-continuous and increasing function. A function $f \in H(\mathbb{D})$ belongs to the space \mathcal{Q}_K if

$$\|f\|_{\mathcal{Q}_K}^2 = \sup_{a \in \mathbb{D}} \int_{\mathbb{D}} |f'(z)|^2 K(g(a, z)) dA(z) < \infty,$$

where dA is the area measure on \mathbb{D} and $g(a, z)$ is the Green function in \mathbb{D} with singularity at $a \in \mathbb{D}$. By [5, Theorem 2.1], we know that $\|f\|_{\mathcal{Q}_K}^2$ is equivalent to

$$\sup_{a \in \mathbb{D}} \int_{\mathbb{D}} |f'(z)|^2 K(1 - |\sigma_a(z)|^2) dA(z),$$

where $\sigma_a(z) = \frac{a-z}{1-\bar{a}z}$ is a Möbius transformation of \mathbb{D} . If $K(t) = t^p$, $0 \leq p < \infty$, then the space \mathcal{Q}_K gives the space \mathcal{Q}_p (cf. [12, 13]). In particular, \mathcal{Q}_0 is the Dirichlet space; $\mathcal{Q}_1 = BMOA$, the space of functions with bounded mean oscillation on \mathbb{D} ; \mathcal{Q}_p is the Bloch space for all $p > 1$. See [5] and [6] for more results on \mathcal{Q}_K spaces. Let $\mathcal{Q}_K(\partial\mathbb{D})$ be the space of $f \in L^2(\partial\mathbb{D})$ with

$$\|f\|_{\mathcal{Q}_K(\partial\mathbb{D})}^2 = \sup_{I \subset \partial\mathbb{D}} \int_I \int_I \frac{|f(\zeta) - f(\eta)|^2}{|\zeta - \eta|^2} K\left(\frac{|\zeta - \eta|}{|I|}\right) |d\zeta| |d\eta| < \infty.$$

Clearly, if $K(t) = t^2$, then $\mathcal{Q}_K(\partial\mathbb{D})$ is equal to $BMO(\partial\mathbb{D})$, the space of functions having bounded mean oscillation on $\partial\mathbb{D}$ (see [7]).

To study \mathcal{Q}_K and $\mathcal{Q}_K(\partial\mathbb{D})$, we usually need two constraints on K as follows.

$$(1.1) \quad \int_0^1 \frac{\varphi_K(s)}{s} ds < \infty$$

1991 *Mathematics Subject Classification.* Primary 30D50, 30H25, 46E15.

Key words and phrases. \mathcal{Q}_K spaces; absolute values; inner-outer factorisation.

The work is supported by NSF of China (No. 11171203, No. 11371234 and No. 11526131), NSF of Guangdong Province (No. 2014A030313471) and Project of International Science and Technology Cooperation Innovation Platform in Universities in Guangdong Province (No. 2014KGJHZ007).

and

$$(1.2) \quad \int_1^\infty \frac{\varphi_K(s)}{s^2} ds < \infty,$$

where

$$\varphi_K(s) = \sup_{0 < t \leq 1} K(st)/K(t), \quad 0 < s < \infty.$$

If K satisfies (1.2), then $\mathcal{Q}_K \subsetneq BMOA \subsetneq H^2$, where H^2 denotes the Hardy space in \mathbb{D} (see [4, 7]). Thus, if K satisfies (1.2), then the function $f \in \mathcal{Q}_K$ has its non-tangential limit \tilde{f} almost everywhere on $\partial\mathbb{D}$. Using the triangle inequality, one gets that if $g \in \mathcal{Q}_K(\partial\mathbb{D})$, then $|g|$ also belongs to $\mathcal{Q}_K(\partial\mathbb{D})$. In general, the converse is not true. Consider

$$g(e^{it}) = \begin{cases} \log t, & 0 < t < \pi, \\ -\log |t|, & -\pi < t < 0. \end{cases}$$

By [8, p. 66], $|g| \in BMO(\partial\mathbb{D})$, but $g \notin BMO(\partial\mathbb{D})$. For $g \in H^2$, it is natural to seek a condition together with $|\tilde{g}| \in \mathcal{Q}_K(\partial\mathbb{D})$ is equivalent to $g \in \mathcal{Q}_K$. Our main result, (iii) of Theorem 1.1, is even new for \mathcal{Q}_p spaces.

Theorem 1.1. *Suppose that K satisfies (1.1) and (1.2). Let $f \in H^2$. Set*

$$d\mu_z(\zeta) = \frac{1 - |z|^2}{2\pi|\zeta - z|^2} |d\zeta|, \quad z \in \mathbb{D}, \quad \zeta \in \partial\mathbb{D}.$$

Then the following conditions are equivalent.

- (i) $\tilde{f} \in \mathcal{Q}_K$.
- (ii) $\tilde{f} \in \mathcal{Q}_K(\partial\mathbb{D})$.
- (iii) $|f| \in \mathcal{Q}_K(\partial\mathbb{D})$ and

$$(1.3) \quad \sup_{a \in \mathbb{D}} \int_{\mathbb{D}} \left(\int_{\partial\mathbb{D}} |\tilde{f}(\zeta)| d\mu_z(\zeta) - |f(z)| \right)^2 \frac{K(1 - |\sigma_a(z)|^2)}{(1 - |z|^2)^2} dA(z) < \infty.$$

Applying Theorem 1.1, in Section 4, we will show a new criterion for inner-outer factorisation of \mathcal{Q}_K spaces.

In this article, the symbol $A \approx B$ means that $A \lesssim B \lesssim A$. We say that $A \lesssim B$ if there exists a constant C such that $A \leq CB$.

2. PRELIMINARIES

Given $f \in L^2(\partial\mathbb{D})$, let \hat{f} be the Poisson extension of f . Namely,

$$\hat{f}(z) = \int_{\partial\mathbb{D}} f(\zeta) d\mu_z(\zeta), \quad z \in \mathbb{D}.$$

We first give the following characterization of $\mathcal{Q}_K(\partial\mathbb{D})$ spaces. In particular, if $K(t) = t^p$, $0 < p < 1$, the corresponding result was proved in [11].

Theorem 2.1. *Suppose that K satisfies (1.1) and (1.2). Let $f \in L^2(\partial\mathbb{D})$. Then $f \in \mathcal{Q}_K(\partial\mathbb{D})$ if and only if*

$$(2.1) \quad \sup_{a \in \mathbb{D}} \int_{\mathbb{D}} \left(\int_{\partial\mathbb{D}} |f(\zeta)|^2 d\mu_z(\zeta) - |\widehat{f}(z)|^2 \right) \frac{K(1 - |\sigma_a(z)|^2)}{(1 - |z|^2)^2} dA(z) < \infty.$$

To prove Theorem 2.1, we need the following estimate.

Lemma 2.2. *Let (1.1) and (1.2) hold for K . If $s < 1 + c$ and $2s + r - 4 \geq 0$, then*

$$\int_{\mathbb{D}} \frac{K(1 - |\sigma_a(w)|^2)}{(1 - |w|^2)^s |1 - \overline{w}z|^r} dA(w) \approx \frac{K(1 - |\sigma_a(z)|^2)}{(1 - |z|^2)^{s+r-2}}$$

for all $a, z \in \mathbb{D}$. Here c is a small enough positive constant which depends only on (1.1) and (1.2).

Proof. We point out that

$$\int_{\mathbb{D}} \frac{K(1 - |\sigma_a(w)|^2)}{(1 - |w|^2)^s |1 - \overline{w}z|^r} dA(w) \lesssim \frac{K(1 - |\sigma_a(z)|^2)}{(1 - |z|^2)^{s+r-2}}$$

was proved in [1]. So we need only to prove the reverse. For any $z \in \mathbb{D}$, let

$$E(z, 1/2) = \{w \in \mathbb{D} : |\sigma_z(w)| < 1/2\}$$

be the pseudo-hyperbolic disk. It is well known that

$$1 - |z| \approx 1 - |w| \approx |1 - \overline{w}z|$$

for all $w \in E(z, 1/2)$. Furthermore, by [14, Lemma 4.30], we have that $|1 - a\overline{w}| \approx |1 - a\overline{z}|$ for all $a \in \mathbb{D}$ and $w \in E(z, 1/2)$. Since K satisfies (1.2), $K(2t) \approx K(t)$ for all $t \in (0, 1)$. We obtain

$$\begin{aligned} \int_{\mathbb{D}} \frac{K(1 - |\sigma_a(w)|^2)}{(1 - |w|^2)^s |1 - \overline{w}z|^r} dA(w) &\geq \int_{E(z, 1/2)} \frac{K(1 - |\sigma_a(w)|^2)}{(1 - |w|^2)^s |1 - \overline{w}z|^r} dA(w) \\ &\approx \frac{K(1 - |\sigma_a(z)|^2)}{(1 - |z|^2)^{s+r-2}}, \end{aligned}$$

which gives the desired result. \square

Proof of Theorem 2.1. For any $f \in L^2(\partial\mathbb{D})$, the Littlewood-Paley identity ([7, p. 228]) shows that

$$(2.2) \quad \int_{\mathbb{D}} |\nabla \widehat{f}(w)|^2 \log \frac{1}{|w|} dA(w) = \frac{1}{2\pi} \int_{\partial\mathbb{D}} |f(\zeta) - \widehat{f}(0)|^2 |d\zeta|.$$

Replacing \widehat{f} by $\widehat{f \circ \sigma_z}$ in (2.2) for $z \in \mathbb{D}$, one obtains

$$\int_{\partial\mathbb{D}} |f(\zeta)|^2 d\mu_z(\zeta) - |\widehat{f}(z)|^2 \approx \int_{\mathbb{D}} |\nabla \widehat{f}(w)|^2 (1 - |\sigma_z(w)|^2) dA(w).$$

Using Fubini's theorem and Lemma 2.2, we obtain, for all $a \in \mathbb{D}$, that

$$\int_{\mathbb{D}} \left(\int_{\partial\mathbb{D}} |f(\zeta)|^2 d\mu_z(\zeta) - |\widehat{f}(z)|^2 \right) \frac{K(1 - |\sigma_a(z)|^2)}{(1 - |z|^2)^2} dA(z)$$

$$\begin{aligned}
&\approx \int_{\mathbb{D}} \left(\int_{\mathbb{D}} |\nabla \widehat{f}(w)|^2 (1 - |\sigma_z(w)|^2) dA(w) \right) \frac{K(1 - |\sigma_a(z)|^2)}{(1 - |z|^2)^2} dA(z) \\
&\approx \int_{\mathbb{D}} |\nabla \widehat{f}(w)|^2 dA(w) \int_{\mathbb{D}} \frac{(1 - |\sigma_z(w)|^2) K(1 - |\sigma_a(z)|^2)}{(1 - |z|^2)^2} dA(z) \\
&\approx \int_{\mathbb{D}} |\nabla \widehat{f}(w)|^2 K(1 - |\sigma_a(w)|^2) dA(w).
\end{aligned}$$

By [9], we know that $f \in \mathcal{Q}_K(\partial\mathbb{D})$ if and only if

$$\sup_{a \in \mathbb{D}} \int_{\mathbb{D}} |\nabla \widehat{f}(z)|^2 K(1 - |\sigma_a(z)|^2) dA(z) < \infty.$$

Therefore, $f \in \mathcal{Q}_K(\partial\mathbb{D})$ if and only if

$$\sup_{a \in \mathbb{D}} \int_{\mathbb{D}} \left(\int_{\partial\mathbb{D}} |f(\zeta)|^2 d\mu_z(\zeta) - |\widehat{f}(z)|^2 \right) \frac{K(1 - |\sigma_a(z)|^2)}{(1 - |z|^2)^2} dA(z) < \infty.$$

□

By [6], if (1.1) and (1.2) hold for K and $f \in H^2$, then $f \in \mathcal{Q}_K$ if and only if $\widetilde{f} \in \mathcal{Q}_K(\partial\mathbb{D})$. This together with Theorem 2.1, gives the following result immediately which was also obtained in [10] by a different method.

Corollary 2.3. *Suppose that K satisfies (1.1) and (1.2). Let $f \in H^2$. Then $f \in \mathcal{Q}_K$ if and only if*

$$\sup_{a \in \mathbb{D}} \int_{\mathbb{D}} \left(\int_{\partial\mathbb{D}} |\widetilde{f}(\zeta)|^2 d\mu_z(\zeta) - |f(z)|^2 \right) \frac{K(1 - |\sigma_a(z)|^2)}{(1 - |z|^2)^2} dA(z) < \infty.$$

3. PROOF OF THEOREM 1.1

Recall that $B \in H(\mathbb{D})$ is called an inner function if B is bounded in \mathbb{D} and $|\widetilde{B}(\zeta)| = 1$ for almost every $\zeta \in \partial\mathbb{D}$. An outer function for the Hardy space H^2 is the function of the form

$$O(z) = \eta \exp \left(\int_{\partial\mathbb{D}} \frac{\zeta + z}{\zeta - z} \log \psi(\zeta) \frac{d\zeta}{2\pi} \right), \quad \eta \in \partial\mathbb{D},$$

where $\psi > 0$ a.e. on $\partial\mathbb{D}$, $\log \psi \in L^1(\partial\mathbb{D})$ and $\psi \in L^2(\partial\mathbb{D})$. See [4] for more results on inner and outer functions. Using a technique in [2], we give the proof of Theorem 1.1 as follows.

Proof of Theorem 1.1. Note that (i) \Leftrightarrow (ii) was proved in [6].

(i) \Rightarrow (iii). For $f \in \mathcal{Q}_K$, we have $\widetilde{f} \in \mathcal{Q}_K(\partial\mathbb{D})$. The triangle inequality gives that $|\widetilde{f}| \in \mathcal{Q}_K(\partial\mathbb{D})$. For any $z \in \mathbb{D}$, it follows by Hölder's inequality that

$$\begin{aligned}
\left(\int_{\partial\mathbb{D}} |\widetilde{f}(\zeta)| d\mu_z(\zeta) - |f(z)| \right)^2 &\leq \left(\int_{\partial\mathbb{D}} |\widetilde{f}(\zeta) - f(z)| d\mu_z(\zeta) \right)^2 \\
&\leq \int_{\partial\mathbb{D}} |\widetilde{f}(\zeta) - f(z)|^2 d\mu_z(\zeta) \\
&= \int_{\partial\mathbb{D}} |\widetilde{f}(\zeta)|^2 d\mu_z(\zeta) - |f(z)|^2.
\end{aligned}$$

Since $f \in \mathcal{Q}_K$, the above estimate, together with Corollary 2.3, gives (1.3).

(iii) \Rightarrow (i). If $f \equiv 0$, the result is true. Note that $f \in H^2$. If $f \not\equiv 0$, then f must be of the form BO , where B is an inner function and O is an outer function of H^2 (see [4]). By the estimates of B and O respectively, Böe [2, p. 237] gave that for any $z \in \mathbb{D}$,

$$|f'(z)| \leq \frac{4}{1-|z|} \left(\int_{\partial\mathbb{D}} \left| |\tilde{f}(\zeta)| - |\widehat{f}|(z) \right| d\mu_z(\zeta) + |\widehat{f}|(z) - |f(z)| \right).$$

Here we remind that

$$|\widehat{f}|(z) = \int_{\partial\mathbb{D}} |\tilde{f}(\zeta)| d\mu_z(\zeta).$$

Thus, for any $a \in \mathbb{D}$, by Hölder's inequality, we deduce that

$$\begin{aligned} & \int_{\mathbb{D}} |f'(z)|^2 K(1-|\sigma_a(z)|) dA(z) \\ & \lesssim \int_{\mathbb{D}} \left(\int_{\partial\mathbb{D}} \left| |\tilde{f}(\zeta)| - |\widehat{f}|(z) \right| d\mu_z(\zeta) \right)^2 \frac{K(1-|\sigma_a(z)|)}{(1-|z|^2)^2} dA(z) \\ & \quad + \int_{\mathbb{D}} \left(|\widehat{f}|(z) - |f(z)| \right)^2 \frac{K(1-|\sigma_a(z)|)}{(1-|z|^2)^2} dA(z) \\ & \lesssim \int_{\mathbb{D}} \left(\int_{\partial\mathbb{D}} \left(|\tilde{f}(\zeta)| - |\widehat{f}|(z) \right)^2 d\mu_z(\zeta) \right) \frac{K(1-|\sigma_a(z)|)}{(1-|z|^2)^2} dA(z) \\ & \quad + \int_{\mathbb{D}} \left(|\widehat{f}|(z) - |f(z)| \right)^2 \frac{K(1-|\sigma_a(z)|)}{(1-|z|^2)^2} dA(z) \\ & \approx \int_{\mathbb{D}} \left(\int_{\partial\mathbb{D}} |\tilde{f}(\zeta)|^2 d\mu_z(\zeta) - \left(|\widehat{f}|(z) \right)^2 \right) \frac{K(1-|\sigma_a(z)|)}{(1-|z|^2)^2} dA(z) \\ & \quad + \int_{\mathbb{D}} \left(|\widehat{f}|(z) - |f(z)| \right)^2 \frac{K(1-|\sigma_a(z)|)}{(1-|z|^2)^2} dA(z). \end{aligned}$$

By Theorem 2.1 and (1.3), $f \in \mathcal{Q}_K$. The proof is complete. \square

Remark. J. Xiao [11] gave an interesting characterization of \mathcal{Q}_p spaces in terms of functions with absolute values. Namely, if $0 < p < 1$ and $f \in H^2$, then $f \in \mathcal{Q}_p$ if and only if $|\tilde{f}| \in \mathcal{Q}_p(\partial\mathbb{D})$ and

$$\sup_{a \in \mathbb{D}} \int_{\mathbb{D}} \left(\left(\int_{\partial\mathbb{D}} |\tilde{f}(\zeta)| d\mu_z(\zeta) \right)^2 - |f(z)|^2 \right) \frac{(1-|\sigma_a(z)|^2)^p}{(1-|z|^2)^2} dA(z) < \infty.$$

Our Theorem 1.1 implies Xiao's result above. In fact, set $K(t) = t^p$, $0 < p < 1$, in our Theorem 1.1 and Corollary 2.3. Note that

$$\left(\int_{\partial\mathbb{D}} |\tilde{f}(\zeta)| d\mu_z(\zeta) \right)^2 - |f(z)|^2 \geq \left(\int_{\partial\mathbb{D}} |\tilde{f}(\zeta)| d\mu_z(\zeta) - |f(z)| \right)^2$$

and

$$\left(\int_{\partial\mathbb{D}} |\tilde{f}(\zeta)| d\mu_z(\zeta) \right)^2 \leq \int_{\partial\mathbb{D}} |\tilde{f}(\zeta)|^2 d\mu_z(\zeta).$$

Thus, one can obtain Xiao's result directly.

4. AN APPLICATION TO INNER-OUTER FACTORISATION OF \mathcal{Q}_K SPACES

In this section, we will show a new criterion for inner-outer decomposition of \mathcal{Q}_K spaces. In fact, an inner-outer factorisation characterization of \mathcal{Q}_K spaces has been obtained in [6] as follows.

Theorem A. *Let K satisfy (1.1) and (1.2) with*

$$\tilde{K}(|z|^2) = -\frac{\partial^2 K(1 - |z|^2)}{\partial z \partial \bar{z}}, \quad z \in \mathbb{D}.$$

Let $f \in H^2$ with $f \not\equiv 0$. Then $f \in \mathcal{Q}_K$ if and only if $f = BO$, where B is an inner function and O is an outer function in \mathcal{Q}_K for which

$$(4.1) \quad \sup_{a \in \mathbb{D}} \int_{\mathbb{D}} |O(z)|^2 (1 - |B(z)|^2) \tilde{K}(|\sigma_a(z)|^2) |\sigma'_a(z)|^2 dA(z) < \infty.$$

As an application of Theorem 1.1, we obtain the following result.

Theorem 4.1. *Let K satisfy (1.1) and (1.2) with*

$$\tilde{K}(|z|^2) = -\frac{\partial^2 K(1 - |z|^2)}{\partial z \partial \bar{z}}, \quad z \in \mathbb{D}.$$

Let $f \in H^2$ with $f \not\equiv 0$. Then $f \in \mathcal{Q}_K$ if and only if $f = BO$, where B is an inner function and O is an outer function in \mathcal{Q}_K for which

$$(4.2) \quad \sup_{a \in \mathbb{D}} \int_{\mathbb{D}} |O(z)|^2 (1 - |B(z)|^2)^2 \tilde{K}(|\sigma_a(z)|^2) |\sigma'_a(z)|^2 dA(z) < \infty.$$

Remark. Theorem 4.1 shows that formula (4.1) in Theorem A can be replaced by the weaker condition (4.2), and this result is also new for \mathcal{Q}_p spaces.

Proof. Necessity. This is a direct result from Theorem A.

Sufficiency. Let $f = BO$ and $O \in \mathcal{Q}_K$. Note that $O \in \mathcal{Q}_K$ is equivalent to $\tilde{O} \in \mathcal{Q}_K(\partial\mathbb{D})$. By the triangle inequality, one gets $|\tilde{O}| \in \mathcal{Q}_K(\partial\mathbb{D})$. Hence $|\tilde{f}| \in \mathcal{Q}_K(\partial\mathbb{D})$. Observe that

$$(4.3) \quad \int_{\partial\mathbb{D}} |\tilde{f}(\zeta)| d\mu_z(\zeta) - |f(z)| = \int_{\partial\mathbb{D}} |\tilde{O}(\zeta)| d\mu_z(\zeta) - |O(z)| + |O(z)| - |B(z)O(z)|.$$

Wulan and Ye [10] gave that if K satisfies (1.1) and (1.2), then for all $z \in \mathbb{D}$

$$(4.4) \quad \tilde{K}(|z|^2) \approx \frac{K(1 - |z|^2)}{(1 - |z|^2)^2}.$$

By Hölder's inequality, $\tilde{O} \in \mathcal{Q}_K(\partial\mathbb{D})$ and Corollary 2.3, we show that for any $a \in \mathbb{D}$,

$$\begin{aligned} & \int_{\mathbb{D}} \left(\int_{\partial\mathbb{D}} |\tilde{O}(\zeta)| d\mu_z(\zeta) - |O(z)| \right)^2 \frac{K(1 - |\sigma_a(z)|^2)}{(1 - |z|^2)^2} dA(z) \\ & \leq \int_{\mathbb{D}} \left(\int_{\partial\mathbb{D}} |\tilde{O}(\zeta) - O(z)|^2 d\mu_z(\zeta) \right) \frac{K(1 - |\sigma_a(z)|^2)}{(1 - |z|^2)^2} dA(z) \\ & = \int_{\mathbb{D}} \left(\int_{\partial\mathbb{D}} |\tilde{O}(\zeta)|^2 d\mu_z(\zeta) - |O(z)|^2 \right) \frac{K(1 - |\sigma_a(z)|^2)}{(1 - |z|^2)^2} dA(z) < \infty. \end{aligned}$$

Combining the above inequality, (4.2), (4.3) and (4.4), we get

$$\sup_{a \in \mathbb{D}} \int_{\mathbb{D}} \left(\int_{\partial \mathbb{D}} |\tilde{f}(\zeta)| d\mu_z(\zeta) - |f(z)| \right)^2 \frac{K(1 - |\sigma_a(z)|^2)}{(1 - |z|^2)^2} dA(z) < \infty.$$

Applying Theorem 1.1, we get $f \in \mathcal{Q}_K$. The proof is complete. \square

For $f \in \mathcal{Q}_K \subseteq H^2$, if we ignore the choice of a constant with modulus one, then f has a unique decomposition with the form $f(z) = B(z)O(z)$, where B is an inner function and O is an outer function. Combining this with Theorem A and Theorem 4.1, we obtain an interesting result as follows.

Corollary 4.2. *Suppose that K satisfies (1.1) and (1.2). Let B be an inner function and let O be an outer function in \mathcal{Q}_K . Then the following conditions are equivalent.*

(i) *For some $p \in [1, 2]$,*

$$\sup_{a \in \mathbb{D}} \int_{\mathbb{D}} |O(z)|^2 (1 - |B(z)|^2)^p \tilde{K}(|\sigma_a(z)|^2) |\sigma'_a(z)|^2 dA(z) < \infty.$$

(ii) *For all $p \in [1, 2]$,*

$$\sup_{a \in \mathbb{D}} \int_{\mathbb{D}} |O(z)|^2 (1 - |B(z)|^2)^p \tilde{K}(|\sigma_a(z)|^2) |\sigma'_a(z)|^2 dA(z) < \infty.$$

REFERENCES

- [1] G. Bao, Z. Lou, R. Qian and H. Wulan, *Improving multipliers and zero sets in \mathcal{Q}_K spaces*, Collect. Math. **66** (2015), no. 3, 453-468.
- [2] B. Bøe, *A norm on the holomorphic Besov space*, Proc. Amer. Math. Soc. **131** (2003), no. 1, 235-241.
- [3] K. Dyakonov, *Besov spaces and outer functions*, Michigan Math. J. **45** (1998), no. 1, 143-157.
- [4] P. Duren, *Theory of H^p Spaces*, Academic Press, New York, 1970.
- [5] M. Essén and H. Wulan, *On analytic and meromorphic function and spaces of \mathcal{Q}_K -type*, Illinois. J. Math. **46** (2002), no. 4, 1233-1258.
- [6] M. Essén, H. Wulan and J. Xiao, *Several function-theoretic characterizations of Möbius invariant \mathcal{Q}_K spaces*, J. Funct. Anal. **230** (2006), no. 1, 78-115.
- [7] J. Garnett, *Bounded Analytic Functions*, Springer, New York, 2007.
- [8] D. Girela, *Analytic functions of bounded mean oscillation*, In: Complex Function Spaces, Mekrijärvi 1999 Editor: R. Aulaskari. Univ. Joensuu Dept. Math. Rep. Ser. 4, Univ. Joensuu, Joensuu (2001) pp. 61-170.
- [9] J. Pau, *Bounded Möbius invariant \mathcal{Q}_K spaces*, J. Math. Anal. Appl. **338** (2008), no. 2, 1029-1042.
- [10] H. Wulan and F. Ye, *Some results in Möbius invariant \mathcal{Q}_K spaces*, Complex Var. Elliptic Equ. **60** (2015), no. 11, 1602-1611.
- [11] J. Xiao, *Some results on \mathcal{Q}_p spaces, $0 < p < 1$, continued*, Forum Math. **17** (2005), no. 4, 637-668.
- [12] J. Xiao, *Holomorphic \mathcal{Q} Classes*, Springer, LNM 1767, Berlin, 2001.
- [13] J. Xiao, *Geometric \mathcal{Q}_p Functions*, Birkhäuser Verlag, Basel-Boston-Berlin, 2006.
- [14] K. Zhu, *Operator Theory in Function Spaces*, American Mathematical Society, Providence, RI, 2007.

GUANLONG BAO, DEPARTMENT OF MATHEMATICS, SHANTOU UNIVERSITY, SHANTOU, GUANGDONG 515063, CHINA

E-mail address: glbaoah@163.com

ZENGJIAN LOU, DEPARTMENT OF MATHEMATICS, SHANTOU UNIVERSITY, SHANTOU, GUANGDONG 515063, CHINA

E-mail address: zjlou@stu.edu.cn

RUISHEN QIAN, SCHOOL OF MATHEMATICS AND COMPUTATION SCIENCE, LINGNAN NORMAL UNIVERSITY, ZHANJIANG, GUANGDONG 524048, CHINA

E-mail address: qianruishen@sina.cn

HASI WULAN, DEPARTMENT OF MATHEMATICS, SHANTOU UNIVERSITY, SHANTOU, GUANGDONG 515063, CHINA

E-mail address: wulan@stu.edu.cn